

RELATIVISTIC FREQUENCY IN TRIGONOMETRIC TERMS.

S. Kanagaraj
Euclidean Relativity
s.kana.raj@gmail.com
(24 October 2009)

Abstract

The frequency variation equation is derived in relation to 4-Euclidean metric diagram. The relativistic equations are expressed in terms of a real angular rotation ϕ (or alternatively θ) in trigonometric form. The very small relativistic effect for sound waves due to time dilation is also shown.

1.0 Introduction.

Based on a Euclidean interpretation of special relativity (SR), the Euclidean space-time (EST) diagram was formulated[1] and subsequently transformed into a Euclidean space (ES) analogue of a moving body[2]. The relativistic space, time, momentum and energy variation equations were derived expressed in trigonometric form as a function of the spacetime angle θ (or alternatively the orientation ϕ). We next express the relativistic frequency as a function of θ (or ϕ) in trigonometric form.

2.0 Effects contributing to frequency variation.

When a wavesource with rest frequency ν_o is receding at velocity v , the effect of its position changes causes the wavelength λ received to be larger than when at rest λ_o . The time interval between wavefronts, period T , is larger than T_o when at rest. For this case, $\lambda > \lambda_o$; $T > T_o$ and since $\nu = \frac{1}{T}$, $\nu < \nu_o$. For approaching case $\lambda < \lambda_o$; $T < T_o$ and $\nu > \nu_o$. The wavelength change,

$\Delta\lambda$, with λ_o ratio due to this effect is¹ $\frac{\Delta\lambda}{\lambda_o} = \frac{v}{c_p}$ with c_p , the wave propagation velocity. Re-

arranging, $\frac{\lambda}{\lambda_o} = 1 \pm \frac{v}{c_p}$, with (+) and (-) signs for receding and approaching motion

respectively. Noting $\nu\lambda = c_p$ and assuming constant propagation velocity, $\frac{\lambda}{\lambda_o} = \frac{\nu_o}{\nu} = \frac{T}{T_o}$.

From EST diagram, $\cos\theta = \frac{v}{c}$, where c is the invariant velocity in spacetime. If light propagation velocity is c_{light} then $c_p = c_{light} = c$ and

¹ This relationship for a wave source moving along a straight path from a stationary observer is derived with ease from first principles.

$$\frac{v_o}{v} = 1 + \cos \theta \quad \dots\dots \text{Eqn 1}$$

When receding, v ranges from 0 to c and the corresponding θ range is $\pi/2 \geq \theta \geq 0$. When approaching, v ranges from 0 to $-c$ and θ range is $\pi/2 \leq \theta \leq \pi$. The $\cos \theta$ sign change is consistent with the velocity sign change for different directions.

Also due to time dilation (TD), a moving wavesource frequency ν is less than when at rest ν_o .

The change in the periods ratio $\frac{T}{T_o} \left(= \frac{\nu_o}{\nu} \right)$ due to this effect is the TD ratio, thus

$$\frac{\nu_o}{\nu} = \frac{T}{T_o} = TD = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}.$$

Substituting $\cos \theta = \frac{v}{c}$, $\frac{\nu_o}{\nu} = \text{cosec } \theta \quad \dots\dots\dots \text{Eqn 2}$

For receding and approaching motion, θ ranges from $\pi/2 \geq \theta \geq 0$ and $\pi/2 \leq \theta \leq \pi$ respectively with $\text{cosec } \theta$ sign consistent with time variation as independent of these directions.

3.0 Deriving the frequency variation equation.

The resultant frequency variation is the product of the two effects in *Eqn 1* and *2*.

For the case of a moving lightwave source,

$$\begin{aligned} \frac{\nu_o}{\nu} &= (1 + \cos \theta) \text{cosec } \theta \\ &= \text{cosec } \theta + \cot \theta \quad \dots\dots \text{Eqn 3} \end{aligned}$$

From *Eqn 3*,

$$\frac{\nu_o}{\nu} = \frac{1 + \cos \theta}{\sin \theta} = \sqrt{\frac{1 + \cos \theta}{1 - \cos \theta}}$$

Since $\cos \theta = \frac{v}{c}$, $\frac{\nu_o}{\nu} = \sqrt{\frac{1 + \frac{v}{c}}{1 - \frac{v}{c}}}$ for receding motion with the signs reversed for approaching

motion, consistent with the relativistic Doppler frequency shift for light. For a soundwave source,

$$\frac{v_o}{v} = \left(1 \pm \frac{v}{c_{sound}} \right) cosec \theta \quad \dots\dots Eqn 4$$

where c_{sound} is the soundwave propagation velocity. As c is the invariant velocity in spacetime, the effect due to time dilation remains as $cosec \theta$. Eqn 4 which accounts for time dilation is consistent with the sound Doppler effect [3]. Since $cosec \theta = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}} = TD$ and for this case

$v \ll c$, $cosec \theta$ is very close to 1. The time dilation correction factor is negligible [4] and reduces to classical Doppler shift equation.

Since $cos \theta = \frac{v}{c} = sin \phi$, substituting into Eqn 3, the equation is expressed alternatively as a function of the orientation angle ϕ with the sign change again consistent with relativistic requirements,

$$\frac{v_o}{v} = sec \phi + tan \phi \quad \dots\dots\dots Eqn 5$$

In conclusion, the derived expressions provides an alternative approach to investigate frequency variations as a function of θ or ϕ in trigonometric terms and offers new avenues in studying relativistic dynamics.

REFERENCES

1. Kanagaraj, S : *Euclidean alternative to Minkowski spacetime diagram.*, <http://www.euclideanrelativity.net> , 12 Aug 2009.
2. Kanagaraj, S : *Euclidean space analogue of a moving body.*, <http://www.euclideanrelativity.net>, 2 Sept 2009
3. Reynolds, Robert : *Doppler effect for sound via classical and relativistic space-time diagrams*, Am.J.Phys.,58(4),390-394,1990.
4. Bachman, R A : *Relativistic acoustic Doppler effect*, Am.J.Phys.,50,816-818,1982.

----- END -----

© Kanagaraj